
Brevetoxin (NSP) Plate, Detailed ELISA Procedure


1. Addition of Standards, Samples

Add 50 uL of the standard solutions or samples into the wells of the test strips according to the working scheme given. We recommend using duplicates or triplicates.




4. Addition of Substrate/Color Solution

Add 100 uL of substrate/color solution to the individual wells successively using a multichannel pipette or a stepping pipette. Cover the wells with parafilm or tape and mix the contents by moving the strip holder in a rapid circular motion on the benchtop. Be careful not to spill contents. Incubate the strips for 30 min at room temperature.




2. Addition of Reconstituted Enyzme Conjugate

Add 50 uL of the reconstituted enzyme conjugate to the individual wells successively using a multi- channel pipette or a stepping pipette. Cover and mix for 30 seconds by rotating on benchtop. Incubate for 60 minutes at room temperature.




5. Addition of Stopping Solution

Add 100 uL of stop solution to the wells in the same sequence as for the substrate solution using a multi- channel pipette or a stepping pipette.




3. Washing of Plates

After incubation, remove the covering and vigorously shake the contents of the wells into a sink. Wash the strips three times with a multi-channel pipette or wash bottle using the diluted 1X washing buffer solution. Please use at least a volume of 250 uL of washing buffer for each well and each washing step. Remaining buffer in the wells should be removed by patting the plate dry on a stack of paper towels.



6. Measurement of Color

Read the absorbance at 450 nm using a microplate ELISA reader. Calculate results.



Eurofins Abraxis

124 Railroad Drive, Warminster, PA 18974 USA Phone: 215-357-3911 Fax: 215-357-5232

www.abraxiskits.com Technical Support:

Support.ET.Warminster@eurofinsUS.com

Ordering:


Info.ET.Warminster@eurofinsUS.com



Brevetoxin (NSP) Plate, Concise ELISA Procedure


1. Addition of Standards, Samples

Add 50 uL of standard solutions or samples.




4. Addition of Substrate/Color Solution

Add 100 uL of substrate/color solution. Incubate 30 minutes at room temperature and away from direct sunlight.




2. Addition of Reconstituted Enyzme Conjugate

Add 50 uL of reconstituted enzyme conjugate. Cover and mix for 30 seconds by rotating on benchtop. Incubate for 60 minutes at room temperature.




5. Addition of Stopping Solution

Add 100 uL of stop solution.




3. Washing of Plates

Wash the plates three times with 250 uL of diluted 1X washing buffer.



6. Measurement of Color

Measure color at 450 nm. Calculate results.



Eurofins Abraxis

124 Railroad Drive, Warminster, PA 18974 USA Phone: 215-357-3911 Fax: 215-357-5232

www.abraxiskits.com Technical Support:

<u>Support.ET.Warminster@eurofinsUS.com</u> Ordering:

Info.ET.Warminster@eurofinsUS.com

